

Spin Chains and Chiral Lattice Fermions

H. B. Thacker

Dept. of Physics, Univ. of Virginia, Charlottesville, VA 22901 USA *

The generalization of Lorentz invariance to solvable two-dimensional lattice fermion models has been formulated in terms of Baxter's corner transfer matrix. In these models, the lattice Hamiltonian and boost operator are given by fermionized nearest-neighbor Heisenberg spin chain operators. The transformation properties of the local lattice fermion operators under a boost provide a natural and precise way of generalizing the chiral structure of a continuum Dirac field to the lattice. The resulting formulation differs from both the Wilson and staggered (Kogut-Susskind) prescriptions. In particular, an axial Q_5 rotation is sitewise local, while the vector charge rotation mixes nearest neighbors on even and odd sublattices.

One of the most interesting developments in the theory of solvable two-dimensional lattice statistics models is Baxter's invention of the "corner transfer matrix" (CTM)[[1]]. In the same sense that an ordinary (row-to-row) transfer matrix can be viewed as an operator which translates by one row of sites on a lattice, the CTM may be defined as an operator which rotates around a central site from a horizontal row (or half-row) of sites to the corresponding vertical row, or vice-versa. Thus, for example, the partition function may be written as the product of four CTM's. From this point of view, it is not surprising that the CTM is associated with a lattice version of Euclidean rotational or Lorentz symmetry. What is surprising is that, in certain integrable models like the 8-Vertex/XYZ-spin chain, the CTM contains a continuously variable anisotropy ("spectral") parameter which plays the role of rapidity or Euclidean rotation angle. The action of the full one-parameter set of CTM's on the Hamiltonian eigenstates is completely analogous to the action of the Lorentz boost operator in a relativistically invariant continuum theory, i.e. it uniformly shifts the rapidity of all the particles (spin-waves) in the state by a continuously variable boost velocity. The implications of the CTM for the space-time symmetry properties of integrable lattice models has been discussed elsewhere [2,3]. Here I want to show how the CTM formalism may be used to investigate the chiral

structure of the lattice fermions obtained from the 8-vertex/XYZ chain. It was shown long ago by Luther ([4]) that, in the scaling limit, this model reduces to the relativistically invariant massive Thirring model, a theory of self-coupled, massive Dirac fermions. Since the chiral structure of the massive Thirring model may be expressed in terms of the Lorentz transformation properties of the Dirac field, the CTM formalism provides a natural and unique generalization of this chiral structure to the lattice. To summarize the main result of this investigation, we find that the XYZ spin chain Hamiltonian constitutes a formulation of lattice Dirac fermions which differs essentially from both the Wilson and Kogut-Susskind (KS) prescriptions. In particular, for both Wilson and KS formalisms, the complex spinor components of the Dirac field remain sitewise local. (In the KS case, different spinor components are spread over neighboring sites, but each site supports complex fermion components.) Thus, the conserved $U(1)$ vector charge $Q = \int \bar{\psi}\gamma_0\psi dx$ of the Dirac field simply generates a phase rotation at each site. By contrast, the lattice prescription that we infer from the CTM analysis of the vertex/spin-chain model leads to a local axial-vector charge but a nonlocal vector charge operator which mixes even and odd nearest neighbor fermion fields. In fact, the conserved vector charge is not built into the lattice spin-chain theory a priori (as in the Wilson and KS formalisms), but only emerges as a consequence of fermion doubling. The nonlocality of the vector charge results from the reduction of the

*UVA-INPP-94-5;hep-lat/9412019

Brillouin zone, as I will outline here. A more extensive discussion of vertex models, spin chains, and chiral lattice fermions will be presented elsewhere.

The general, anisotropic nearest-neighbor Heisenberg chain Hamiltonian may be written $H = \sum H_j$, where

$$H_j = -\frac{1}{2} \{ \sigma_j^x \sigma_{j+1}^x + k \sigma_j^y \sigma_{j+1}^y + \Delta \sigma_j^z \sigma_{j+1}^z \} \quad (1)$$

The boost operator that emerges from the CTM analysis is the first moment of the same density, $K = \sum [j H_j]$. This model can be converted to a lattice fermion theory by a Jordan-Wigner transformation

$$c_j^{x,y} = \sigma_j^{x,y} \left(\prod_{j' < j} \sigma_{j'}^z \right) \quad (2)$$

Here, c^x and c^y are real canonical fermion fields,

$$(c_j^{x,y})^* = c_j^{x,y}, \quad \{c_j^a, c_{j'}^b\} = \delta_{ab} \delta_{jj'} \quad (3)$$

Expressing the Hamiltonian in terms of a single complex fermion at each site, $c_j = (c_j^x + i c_j^y) / \sqrt{2}$, the terms in H can be written as a kinetic energy term, a mass term, and an interaction term, respectively, $H_j = T_j + M_j + V_j$, where

$$T_j = \frac{1}{2} (1+k) (c_j^* c_{j+1} + c_{j+1}^* c_j) \quad (4)$$

$$M_j = \frac{1}{2} (1-k) (c_j^* c_{j+1}^* + c_{j+1} c_j) \quad (5)$$

$$V_j = -\frac{1}{2} \Delta c_j^* c_j c_{j+1}^* c_{j+1} \quad (6)$$

Note that the mass term violates the symmetry under a global phase rotation of c_j at each site. In the continuum limit, this global phase symmetry is associated with the conservation of axial-vector current, $J_5^\mu = \bar{\psi} \gamma_5 \gamma^\mu \psi$. From here on, let us focus on the free-fermion case $\Delta = 0$. For this case, the eigenstates of $H = T + M$ may be written simply in terms of momentum-space fermion operators,

$$a_{x,y}(z) = \sum_j z^j c_j^{x,y} \quad (7)$$

The eigenmode operators of H are given by

$$B(z) = z^{-1} C(z) a_x(z) + i C(z^{-1}) a_y(z) \quad (8)$$

where $C(z) = (1 + kz^2)^{1/2}$. They satisfy $[H, B(z)] = \omega(z) B(z)$, where the single particle energy is $\omega(z) = C(z) C(z^{-1})$. The boost operator, defined as $K = \sum j [T_j + M_j]$ has the fundamental property [2,3,5] of inducing a uniform rapidity shift on the Hamiltonian eigenmodes, as expressed by the commutator

$$[K, B(z)] = z \omega(z) \frac{\partial}{\partial z} B(z). \quad (9)$$

In order to understand the chiral structure of the local lattice fields, we need to look at their behavior under a boost. When expressed in momentum space, the relationship between the transformation of the local fields and that of the eigenmodes discussed above is quite analogous to the continuum case. After some analysis, we find that the transformation of the local field can be expressed in reduced form by defining

$$\psi_{1,2}(z) = (a_x(z) \pm i \sqrt{k} a_y(z)) / \sqrt{2} \quad (10)$$

which satisfy

$$[K, \psi_{1,2}(z)] = z \frac{\partial}{\partial z} [H, \psi_{1,2}(z)] \pm z^{-1} \sqrt{k} \psi_{1,2}(z) \quad (11)$$

This commutator provides a precise generalization of Lorentz transformations to the lattice. By inspecting the energy function $\omega(z)$ we see that there are two scaling regimes as $k \rightarrow 1$, one at $z \approx i$ and one at $z \approx -i$. In the first region, Eq. (11) reduces to precisely the Lorentz transformation of the chiral components of a Dirac field. Thus if we restrict z to the upper half of the complex plane, the lattice fields $\psi_{1,2}(z)$ are the chiral components of the lattice Dirac field (in momentum space). For the second scaling region at $z \approx -i$ (the “doubler” modes) the Lorentz structure is inverted from what it should be if we try to naively extend this identification to the LHP (e.g. by taking $z \rightarrow -z$ the last term in (11) changes sign). The proper way to deal with the doubler modes is to define a reduced Brillouin zone $0 \leq \arg z < \pi$ by using the reflection relation that follows from the reality of the a_x and a_y fermions, namely

$$\psi_{1,2}(z) = \psi_{2,1}^*(z^*) \quad (12)$$

(In the continuum Dirac theory, this represents CP conjugation.) Any operator involving fermion operators in the lower half z -plane should be rewritten in terms of those in the UHP (using the above reflection formula) in order to correctly exhibit its chiral structure. The doubler modes in the LHP, reflected into the reduced Brillouin zone, provide the CP conjugate states relative to those in the UHP. This has interesting implications for the structure of the vector charge on the lattice. For example, if we define the Fourier transform of the local complex fermion, $\psi(z) = \sum z^j c_j$, we must write this local fermion in terms of Dirac fields of indefinite vector charge. In the reduced Brillouin zone $z \in \text{UHP}$, we can write the fermion fields on even and odd sites separately in terms of the chiral fields (taking $k \rightarrow 1$),

$$\psi_{e,o}(z) \equiv \psi(z) \pm \psi(-z) = \psi_1(z) \pm \psi_2^*(-z^*) \quad (13)$$

It may be verified directly that the Hamiltonian reduces to that of a Dirac fermion with chiral components as indicated. (In the continuum limit, this calculation reduces to the original analysis of Luther[4].)

$$T = \frac{1}{2}(1+k) \int_{\text{UHC}} \frac{dz}{i\pi z} (z + z^{-1}) \quad (14)$$

$$[\psi_1^*(z)\psi_1(z) - \psi_2^*(z)\psi_2(z)] \quad (15)$$

$$M = \frac{1}{2}(1-k) \int_{\text{UHC}} \frac{dz}{i\pi z} (z - z^{-1}) \quad (16)$$

$$[\psi_1^*(z)\psi_2(z) - \psi_2^*(z)\psi_1(z)] \quad (17)$$

where the integrals are over the upper half of the unit circle. Note in particular that with this reduction of the Brillouin zone, the Hamiltonian (including the mass term) exhibits a phase invariance under a vector charge rotation, $\psi_{1,2} \rightarrow e^{i\phi}\psi_{1,2}$. This rotation mixes the local even- and odd-site fermions,

$$\psi_e \rightarrow \cos \phi \psi_e + i \sin \phi \psi_o \quad (18)$$

$$\psi_o \rightarrow \cos \phi \psi_o + i \sin \phi \psi_e \quad (19)$$

Note however that under an axial-vector rotation, $\psi_1 \rightarrow e^{i\phi}\psi_1$, $\psi_2 \rightarrow e^{-i\phi}\psi_2$, the site fermions carry a definite chirality: $\psi_{e,o} \rightarrow e^{i\phi}\psi_{e,o}$.

To understand the local conservation of the vector current in space-time, it is useful to con-

sider the two-dimensional framework of the 8-vertex model and its relation to the Hamiltonian formalism. Each term H_j in the Hamiltonian comes from the expansion of a local vertex connecting sites j and $j+1$. Denoting the fermion states on these two sites by $+$ (occupied) and $-$ (unoccupied), and using (4) and (5), we can write (for $\Delta = 0, k \approx 1$)

$$H_j|++\rangle = \epsilon|--\rangle, \quad H_j|--\rangle = \epsilon|++\rangle \quad (20)$$

$$H_j|+-\rangle = |--\rangle, \quad H_j|-+\rangle = |+-\rangle \quad (21)$$

where $\epsilon = (1-k)/2$ is proportional to the fermion mass. If j is even, the prescription (13) for the Dirac field corresponds to taking a new basis of states denoted by charge numbers associated with densities $\psi_1^*\psi_1$ and $\psi_2^*\psi_2$,

$$|00\rangle = (|+-\rangle - |--\rangle)/\sqrt{2} \quad (22)$$

$$|11\rangle = (|+-\rangle + |--\rangle)/\sqrt{2} \quad (23)$$

$$|01\rangle = |--\rangle, \quad |10\rangle = |++\rangle \quad (24)$$

for which we have

$$H_j|00\rangle = -|00\rangle, \quad H_j|01\rangle = \epsilon|10\rangle \quad (25)$$

$$H_j|10\rangle = \epsilon|01\rangle, \quad H_j|11\rangle = |11\rangle \quad (26)$$

In this second basis, the vector charge is the sum of the two occupation numbers, and is thus conserved through a vertex.

This work was supported in part by the U.S. Department of Energy under grant No. DE-AS05-89ER40518.

REFERENCES

1. R. J. Baxter, Exactly Solved Models in Statistical Mechanics (Academic Press, London, 1982).
2. H. B. Thacker, Physica 18D, 348 (1986).
3. H. Itoyama and H. B. Thacker, Phys. Rev. Lett. 14, 1395 (1987); Nuc. Phys. B320, 541 (1989).
4. A. Luther, Phys. Rev. B14, 2153 (1975).
5. K. Sogo and M. Wadati, Prog. Theo. Phys. 69, 431 (1983).